

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 May 2001 (17.05.2001)

PCT

(10) International Publication Number
WO 01/34118 A2

(51) International Patent Classification⁷: A61K 9/14, 38/55

(21) International Application Number: PCT/US00/30910

(22) International Filing Date:
10 November 2000 (10.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/439,514 12 November 1999 (12.11.1999) US

(71) Applicant: ABBOTT LABORATORIES [US/US]; Dept.
377, Bldg. AP6D-2, 100 Abbott Park Road, Abbott Park,
IL 60064-6050 (US).

(72) Inventors: FORT, James, J.; 2700 Leaffield Terrace, Mid-
lothian, VA 23113 (US). KRILL, Steven, L.; 44 Mey-
ersville Road, Chatham, NJ 07928 (US). LAW, Devalina;
835C Country Club Drive, Libertyville, IL 60048 (US).

QIU, Yihong; 6118 Honeysuckle Lane, Gurnee, IL 60031
(US). PORTER, William, R.; 20 Manchester Lane, Ver-
non Hills, IL 60061 (US). SCHMITT, Eric, A.; 319 Ever-
green Court, Libertyville, IL 60048 (US).

(74) Agents: CROWLEY, Steven, R. et al.; Dept. 377 Bldg.
AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-
6050 (US).

(81) Designated States (*national*): CA, JP, MX.

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).


Published:

— Without international search report and to be republished
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: SOLID DISPERSION PHARMACEUTICAL FORMULATIONS

Amorphous ABT-538 Dispersions in PEG 8000
Dissolution in 0.1N HCl at 37°C



(57) Abstract: A pharmaceutical composition is disclosed which comprises a solid dispersion of an HIV protease inhibitor in a water soluble carrier, such as PEG, having enhanced bioavailability and improved dissolution properties. The solid dispersion may optionally be encapsulated in hard gelatin capsules, compressed into a tablet, or may be granulated with a pharmaceutically acceptable granulating agent. Also disclosed are methods of making said solid dispersion and methods of treating an HIV infection employing said solid dispersion.



WO 01/34118 A2

SOLID DISPERSION PHARMACEUTICAL FORMULATIONS**Technical Field of the Invention**

5 The instant invention relates to the fields of
pharmaceutical and organic chemistry, and provides novel
solid dispersion pharmaceutical formulations with enhanced
bioavailability.

10

Background of the Invention

One measure of the potential usefulness of an oral
dosage form of a pharmaceutical agent is the
15 bioavailability observed after oral administration of the
dosage form. Various factors can affect the
bioavailability of a drug when administered orally. These
factors include aqueous solubility, drug absorption
throughout the gastrointestinal tract, dosage strength, and
20 first pass effect. Aqueous solubility is one of the most
important of these factors. When a drug has poor aqueous
solubility, attempts are often made to identify salts or

other derivatives of the drug which have improved aqueous solubility. When a salt or other derivative of the drug is identified which has good aqueous solubility, it is generally accepted that an aqueous solution formulation of this salt or derivative will provide the optimum oral bioavailability. The bioavailability of the aqueous oral solution formulation of a drug is then generally used as the standard or ideal bioavailability against which other oral dosage forms are measured.

For a variety of reasons, including patient compliance and taste masking, a solid dosage form, such as a capsule or tablet, is usually preferred over a liquid dosage form. However, oral solid dosage forms of a drug generally provide a lower bioavailability than oral solutions of the drug. One goal of the development of a suitable solid dosage form is to obtain a bioavailability of the drug that is as close as possible to the ideal bioavailability demonstrated by the oral aqueous solution formulation of the drug.

An alternative dosage form is a solid dispersion. The term solid dispersion refers to the dispersion of one or more active ingredients in an inert carrier or matrix at

solid state prepared by the melting (or fusion), solvent, or melting-solvent methods. (Chiou and Riegelman, *Journal of Pharmaceutical Sciences*, 60, 1281 (1971)). The dispersion of a drug or drugs in a solid diluent by mechanical mixing is not included in this category. Solid dispersions may also be called solid-state dispersions.

Retroviral protease inhibiting compounds are useful for inhibiting HIV proteases *in vitro* and *in vivo*, and are useful for inhibiting HIV (human immunodeficiency virus) infections and for treating AIDS (acquired immunodeficiency syndrome). HIV protease inhibiting compounds typically are characterized by having poor oral bioavailability.

Examples of HIV protease inhibiting compounds include

2S,3S,5S)-5-(N-(N-(N-methyl-N-(2-isopropyl-4-

thiazolyl)methyl)amino)carbonyl)L-valinyl)amino-2-(N-(5-thiazolyl)methoxy-carbonyl)-amino)-amino-1,6-diphenyl-3-hydroxyhexane (ritonavir);

(2S, 3S, 5S)-2-(2,6-Dimethylphenoxyacetyl)

amino-3-hydroxy-5-[2S-(1-tetrahydro-pyrimid-2-onyl)-3-methyl butanoyl]-amino-1,6-diphenylhexane (ABT-378);

N-(2(R)-hydroxy-1 (S)-indanyl)-2(R)-phenylmethyl

-4 (S) -hydroxy-5- (1- (4- (3-pyridylmethyl) -2 (S) -N' - (t-butylcarboxamido) -piperazinyl)) -pentaneamide (indinavir);

N-tert-butyl-decahydro-2- [2 (R) -hydroxy-4-phenyl-3 (S) - [[N- (2-quinolylcarbonyl) -L-asparaginyll amino] butyl] - (4aS, 8aS) -iso

5 quinoline-3 (S) -carboxamide (saquinavir);

5 (S) -Boc-amino-4 (S) -hydroxy-6-phenyl-2 (R) -phenylmethylhexanoyl- (L) -Val- (L) -Phe-morpholin-4-ylamide;

1 -Naphthoxyacetyl-beta-methylthio-Ala- (2S, 3S) -3-amino-2-hydroxy-4-butanoyl 1,3-thiazolidine-4-

10 t-butylamide;

5-isoquinolinoxyacetyl-beta-methylthio-Ala- (2S, 3S) -3-amino-2-hydroxy-4-butanoyl-1,3-thiazolidine-4-t-butylamide;

[1S- [1R- (R-), 2S*]] -N¹ [3- [[[(1,1 -

15 dimethylethyl) amino] carbonyl] (2-methylpropyl) amino] -2-hydroxy-1- (phenylmethyl) propyl] -2- [(2-quinolinyllcarbonyl) amino] -butanediamide;

VX-478; DMP-323; DMP-450; AG1343 (nelfinavir);

BMS 186,318; SC-55389a; BILA 1096 BS; and U-140690, or

20 combinations thereof.

While some drugs would be expected to have good solubility in organic solvents, it would not necessarily

follow that oral administration of such a solution would give good bioavailability for the drug.

Polyethylene glycol (PEG) solid dispersion formulations are generally known to improve the dissolution and bioavailability of many compounds. However, Aungst et al. has recently demonstrated that this was unable to improve the bioavailability of an HIV protease inhibitor with a cyclic urea structural backbone, called DMP 323 (Aungst et al., *International Journal of Pharmaceutics*, 10 156, 79 (1997)).

Thus, it would be a significant contribution to the art to provide a solid dispersion pharmaceutical formulation of a retroviral protease inhibitor which is more stable and has enhanced bioavailability.

15

Summary of the Invention

The instant invention provides a stable solid dispersion comprising a retroviral protease inhibitor and
5 PEG having improved bioavailability.

Also provided by the instant invention is a pharmaceutical composition comprising a stable solid dispersion as described above with a pharmaceutically acceptable carrier, diluent, or excipient.

10 Additionally provided by the instant invention is a method for preparing a stable solid dispersion as described above.

The instant invention still further provides a method of treating an HIV infection comprising administering an
15 effective amount of a stable solid dispersion as described above to a mammal in need of such treatment.

Brief Description of the Figures

Figure 1 illustrates the dispersion of amorphous ABT-538 in PEG 8000.

5 Figure 2 illustrates the bioavailability of a dispersion of amorphous ABT-538 in PEG 8000.

Figure 3 illustrates the in vivo-in vitro correlation of ABT-538.

Figure 4 illustrates the dissolution of ABT-378.

10 Figure 5 illustrates the dissolution of nelfinavir.

Detailed Description of the Invention

This invention pertains to the preparation of solid dispersion systems for protease inhibitors with improved dissolution and oral bioavailability.

A solid (molecular) dispersion comprising an HIV protease inhibiting compound may be prepared by dissolving or dispersing the HIV protease inhibiting compound in a sufficient amount of an organic solvent followed by dispersion into a suitable water soluble carrier. Suitable organic solvents include pharmaceutically acceptable solvents such as methanol, ethanol, or other organic solvents in which the protease inhibitor is soluble. Suitable water soluble carriers include polymers such as polyethylene glycol (PEG), pluronics, pentaerythritol, pentaerythritol tetraacetate, polyoxyethylene stearates, poly-ε-caprolactone, and the like.

The organic solvent (preferably ethanol) may then be evaporated away, leaving the drug dispersed/dissolved in the molten matrix, which is then cooled. The solid matrix has the compound finely dispersed (molecular dispersion) in such a way that dissolution of the drug is maximized, thus

improving the bioavailability of a drug exhibiting dissolution rate limited absorption. Ease of manufacturing is also an attribute to this type of formulation. Once the organic solvent is evaporated to yield a solid mass, the mass may be ground, sized, and optionally formulated into an appropriate delivery system. Thus, by improving the dissolution of a poorly water soluble drug, the drug in a suitable carrier may be filled into a gelatin capsule as a solid, or the matrix may potentially be compressed into a tablet.

The delivery system of the present invention results in increased solubility and bioavailability, and improved dissolution rate of the HIV protease inhibiting compound.

Other pharmaceutically-acceptable excipients may be added to the formulation prior to forming the desired final product. Suitable excipients include lactose, starch, magnesium stearate, or other pharmaceutically-acceptable fillers, diluents, lubricants, disintegrants, and the like, that might be needed to prepare a capsule or tablet.

The resulting composition comprising the HIV protease inhibiting compound may be dosed directly for oral administration, diluted into an appropriate vehicle for

oral administration, filled into capsules, or made into tablets for oral administration, or delivered by some other means obvious to those skilled in the art. The composition can be used to improve the oral bioavailability and
5 solubility of said HIV protease inhibiting compound.

Total daily dosing of HIV protease inhibitors may be administered to a human in single or divided doses in amounts, for example, from 0.001 to 1000 mg/kg body weight daily, but more usually 0.1 to 50 mg/kg body weight daily.


10 Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose. It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the age, body weight, general health, sex, diet,
15 time of administration, rate of excretion, drugs administered in combination and the severity of the particular disease undergoing therapy.

ABT-538 (ritonavir) was preferably used as the HIV protease inhibitor in the instant invention. Additionally,
20 two other protease inhibitors, ABT-378 and nelfinavir mesylate, were tested in solid dispersions to demonstrate

the improved dissolution which can be achieved with this system.

One aspect of the instant invention provides a solid dispersion of a compound of formula I

5




A compound of formula I is an HIV protease inhibitor

10 marketed by Abbott Laboratories under the tradename Norvir[®],
with the common name ritonavir [(2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-isopropyl-4-thiazolyl)-methyl)amino)carbonyl)-L-valinyl)amino-2-(N-
15 ((5-thiazolyl)methoxy-carbonyl)-amino)-1,6-diphenyl-3-hydroxyhexane]. This and other compounds as well as
methods for preparing same are disclosed in U.S. Patent

Nos. 5,648,497 and 5,541,206, the disclosures of which are herein incorporated by reference.

Additional HIV protease inhibitors which may be formulated into a solid dispersion include compounds of

5 formula II



10


II

A compound of formula II is known as ABT-378 ((2S,3S,5S)-2-(2,6-dimethylphenoxyacetyl)-amino-3-hydroxy-5-(2S-(1-tetrahydropyrimid-2-onyl)-3-methyl-

15 butanoyl)amino-1,6-diphenylhexane). This and other compounds, as well as methods for preparing same, are

identified in U.S. Patent No. 5,914,332, the disclosure of which is herein incorporated by reference.

A compound of formula III provided hereinbelow is known as nelfinavir mesylate (marketed under the tradename
5 Viracept[®] by Agouron Pharmaceuticals, Inc. in La Jolla, CA), and is another HIV protease inhibitor which may be formulated into a solid dispersion.



10

III

The following Examples are provided to further illustrate the present invention.

EXAMPLES

Example 1

5

Dispersion Preparations

A. Ritonavir (ABT-538) Dispersion Preparation:

10

The samples were prepared by dissolving ABT-538 in a small volume of 200 proof ethanol in a 250 ml round bottom flask. The flask was vortexed and then placed in a water bath maintained at 75 °C. The PEG 8000 was added to the hot alcohol solution with continual swirling until the PEG

15 melted. The flask was then attached to a rotary evaporator, immersed in the water bath (75 °C) under vacuum for 15 minutes to remove the ethanol. After the majority of ethanol had evaporated, the flask was immersed in an ice bath for 15 minutes. The contents of the flask were then

20 vacuum dried at room temperature for 6 hours. The solid was transferred to a crystallization dish and placed under vacuum overnight to remove residual ethanol. The material was then ground and sifted. Particles ranging in size from

149 to 420 μm were used for further studies. The drug loads used for these dispersions were 10, 20 and 30% w/w.

B. ABT-378 Dispersion Preparation:

A 10% dispersion was prepared using an alcoholic solution of ABT-378 (ca. 0.1 g/ml) by the same method as described in section A above.

5

C. Nelfinavir mesylate Dispersion Preparation:

Nelfinavir mesylate is available from Agouron Pharmaceuticals, Inc. under the tradename Virucept[®].

A 10% dispersion was prepared using an alcoholic solution of nelfinavir (ca. 0.035 g/ml) by the same method as described in section A above.

The potency values of all the dispersions as well as the dissolution sample concentrations were determined via HPLC.

15

D. Results:

The *in vitro* dissolution data of the ABT-538 dispersions compared with ABT-538 in 0.1N HCl (shown in Figure 1, n=3 \pm SD unless otherwise indicated) show that the dispersions markedly improved the dissolution rate of the drug. Drug loading decreases the rate of drug release in a rank order. A bioavailability study was conducted in

20

dogs with the above ABT-538 dispersions to elicit the drug load effects *in vivo*. Eight beagle dogs, obtained from Marshall Research Animals (North Rose, NY), were utilized in this study. The animals were fasted overnight prior to dosing in each period but water was allowed *ad libitum*. Approximately 30 minutes prior to dosing, each dog received a 100 µg/kg subcutaneous dose of histamine. Capsules containing 5 mg/kg of 10, 20 and 30% solid dispersion (formulations A, B and C, respectively) were tested against crystalline drug as a reference in a four-way crossover study.

Each dog received the dose followed by approximately 10 ml of water. A washout period of approximately 1 week was used to separate each dosing period. The plasma samples were analyzed by a method reported by Marsh *et al.* (Marsh, K.C., Eiden, E. and McDonald, E. Determination of Ritonavir, a new HIV Protease Inhibitor, in Biological Samples Using Reversed-Phase High-Performance Liquid Chromatography. *J. Chromatography B.* 704 (1997) 307-313.)

The results of the study are shown in Figure 2. The results show that the solid dispersions improve absorption compared to the reference. An *in vitro* - *in vivo*

correlation was established. A plot of the AUC versus the amount dissolved in 20 min, shown in Figure 3, is a straight line, indicating excellent correlation.

The dissolution properties of the two additional
5 protease inhibitors (ABT-378 and nelfinavir mesylate) were also determined. The *in vitro* dissolution data (Figure 4) of the ABT-378 dispersion compared with reference clearly shows that the preparation of a dispersion markedly improves dissolution rate of the drug. The variability in
10 the release rate from the dispersion is due to the fact that the preparation of these dispersions had not been optimized to completely overcome the wetting problem of the drug. Despite this, the improvements observed are significant [95% confidence intervals shown].

15 The nelfinavir mesylate solid dispersion also exhibits an improved *in vitro* dissolution rate compared to the neat drug (Figure 5).

E. Conclusions:

20 Solid dispersions of HIV protease inhibitors (for example, ABT-538 (ritonavir), ABT-378, and nelfinavir mesylate) markedly improve the dissolution rate of these

drugs. This improvement of dissolution rate is reflected in the improvement of bioavailability. An excellent *in vivo* - *in vitro* correlation established for the dispersions suggests that the *in vitro* dissolution reflects *in vivo* bioavailability for these systems.

Example 2

Stability of Dispersion in Molten PEG 8000

10

The stability of the dispersion of ABT-538 in PEG 8000 in the molten state at 70 °C was examined. Individual approximately 5 mg quantities of the dispersion (aged for 6 weeks at room temperature) were placed in 4 ml glass vials. These vials, with the exception of the initial time point, were placed in a 70 °C oven which was sampled at pre-determined intervals, chilled in ice water and placed in the freezer until HPLC analysis. After all samples were collected, they were analyzed for ABT-538 content by HPLC. The HPLC system consisted of a Hitachi AS 4000 autosampler, SP 8800 ternary pump, Applied Biosystems 783 detector, and PE Nelson Data

20

acquisition system. Other chromatographic details included a Regis Little Champ 5 cm C-18 column, a mobile phase consisting of an aqueous solution of 0.1% trifluoroacetic acid in 10 mM aqueous tetramethyl ammonium perchlorate (TMAP)/acetonitrile/methanol (55/40/5). The flow rate was 1 ml/minute, the wavelength of detection was 205 nm, and the injection volume was 100 μ l. Standard curves of peak area of ABT-538 vs. concentration in the range of interest were compared with experimentally obtained area counts.

10

Example 3

15 Additional Protocol For Oral Bioavailability Studies

Dogs (beagle dogs, mixed sexes, weighing 7-14 kg) are fasted overnight prior to dosing, but are permitted water *ad libitum*. Each dog receives a 100 μ g/kg subcutaneous dose of histamine approximately 30 minutes prior to dosing. Each dog receives a single solid dosage form corresponding to a 5 mg/kg dose of the drug. The dose is followed by approximately 10 milliliters of water. Blood samples are obtained from each animal prior to dosing and at 0.25, 0.5, 1.0, 1.5, 2, 3, 4, 6, 8, 10 and 12 hours after drug

25

administration. The plasma is separated from the red cells by centrifugation and frozen (- 30 °C) until analysis. The concentrations of parent drug is determined by reverse phase HPLC with low wavelength UV detection following

5 liquid-liquid extraction of the plasma samples. The parent drug area under the curve is calculated by the trapezoidal method over the time course of the study. The absolute bioavailability of each test composition is calculated by comparing the area under the curve after oral dosing to

10 that obtained from a single intravenous dose. Each capsule or capsule composition is evaluated in a group containing at least six dogs. The values reported are averages for each group of dogs.

CLAIMS

What is claimed is:

1. A pharmaceutical composition comprising a solid
5 dispersion of an HIV protease inhibitor or combination of
HIV protease inhibitors and a water soluble carrier.

2. The composition of Claim 1 wherein said water
soluble carrier is polyethylene glycol (PEG).

10

3. The composition of Claim 1 wherein said HIV
protease inhibitor is dissolved in an organic solvent.

4. The composition of Claim 3 wherein said organic
15 solvent is ethanol.

5. The composition of Claim 2 wherein said HIV
protease inhibitor is selected from the group consisting
of:

20 2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-isopropyl-4-
thiazolyl)methyl)amino)carbonyl)L-valinyl)amino-2-(N-(5-

thiazolyl)methoxy-carbonyl)-amino)-amino-1,6-diphenyl-3-hydroxyhexane (ritonavir);

(2S, 3S, 5S)-2-(2,6Dimethylphenoxyacetyl)

amino-3-hydroxy-5-[2S-(1-tetrahydro-pyrimid-2-onyl)-3-methyl butanoyl] amino-1,6-diphenylhexane (ABT-378);

N-(2(R)-hydroxy-1 (S)-indanyl)-2(R)-phenylmethyl

-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide (indinavir);

N-tert-butyl-decahydro-2-[2(R)-hydroxy-4-phenyl-3(S)-[[N-(2-quinolylcarbonyl)-L-asparaginy] amino]butyl]-(4aS,8aS)-isoquinoline-3(S)-carboxamide (saquinavir);

5(S)-Boc-amino-4(S)-hydroxy-6-phenyl-2(R)-

phenylmethylhexanoyl-(L)-Val-(L)-Phe-morpholin-4-ylamide;

1-Naphthoxyacetyl-beta-methylthio-Ala-(2S, 3S)-

3-amino-2-hydroxy-4-butanoyl 1,3-thiazolidine-4-t-butylamide;

5-isoquinolinoxyacetyl-beta-methylthio-Ala-(2S,3S)-3-amino-2-hydroxy-4-butanoyl-1,3-thiazolidine-4-t-butylamide;

[1S-[1R-(R-),2S*]]-N¹-[3-[[[(1,1 -

dimethylethyl) amino] carbonyl] (2-methylpropyl) amino]-2-

hydroxy-1 -(phenylmethyl)propyl]-2-[(2-

quinolinylcarbonyl) amino]-butanediamide;

VX-478; DMP-323; DMP-450; AG1343 (nelfinavir);
BMS 186,318; SC-55389a; BILA 1096 BS; U-140690, or
combinations thereof.

5 6. The composition of Claim 2 wherein said HIV
protease inhibitor is (2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-
isopropyl-4-thiazolyl)methyl)amino)carbonyl)L-
valinyl)amino-2-(N-((5-thiazolyl)methoxy-carbonyl)-amino)-
amino-1,6-diphenyl-3-hydroxyhexane (ritonavir).

10

 7. The composition of Claim 2 wherein said HIV
protease inhibitor is (2S,3S,5S)-2-(2,6-
Dimethylphenoxyacetyl)amino-3-hydroxy-5-[2S-(1-tetrahydro-
pyrimid-2-onyl)-3-methyl-butanoyl] amino-1,6-diphenylhexane
15 (ABT-378).

 8. The composition of Claim 2 wherein said HIV
protease inhibitor is a combination of (2S,3S,5S)-5-(N-(N-
((N-methyl-N-((2-isopropyl-4-
20 thiazolyl)methyl)amino)carbonyl)L-valinyl)amino-2-(N-((5-
thiazolyl)methoxy-carbonyl)-amino)-amino-1,6-diphenyl-3-

hydroxyhexane (ritonavir) and (2S,3S,5S)-2-(2,6-Dimethylphenoxyacetyl)amino-3-hydroxy-5-[2S-(1-tetrahydro-pyrimid-2-onyl)-3-methyl butanoyl] amino-1,6-diphenylhexane (ABT-378).

5

9. The composition of Claim 2 wherein said solid dispersion is encapsulated in a hard gelatin capsule.

10. The composition of Claim 2 wherein said solid
10 dispersion is compressed into a tablet.

11. The composition of Claim 1 further comprising an additive or a mixture of additives independently selected from the group consisting of pharmaceutically acceptable
15 surfactants and antioxidants.

12. A method of preparing a composition of Claim 1 which comprises:

- a) dissolving an HIV protease inhibitor into an
20 organic solvent to form a solution;
- b) adding a water soluble carrier to said solution to form a mixture;

- c) optionally flash evaporating said solvent;
 - d) optionally drying the resulting residue remaining after evaporation;
 - e) optionally grinding and sieving the solid
- 5 dispersion to obtain a resultant product.

13. The method of Claim 12 additionally comprising encapsulating the solid dispersion in a hard gelatin capsule.

10

14. The method of Claim 12 additionally comprising compressing said solid dispersion into a tablet.

15 15. The method of Claim 12 wherein said HIV protease inhibitor is (2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-isopropyl-4-thiazolyl)methyl)amino)carbonyl)L-valinyl)amino-2-(N-((5-thiazolyl)methoxy-carbonyl)-amino)-amino-1,6-diphenyl-3-hydroxyhexane (ritonavir).

20 16. The method of Claim 12 wherein said HIV protease inhibitor is (2S,3S,5S)-2-(2,6-Dimethylphenoxyacetyl)amino-3-hydroxy-5-[2S-(1-tetrahydro-

pyrimid-2-onyl)-3-methyl butanoyl] amino-1,6-diphenylhexane
(ABT-378).


17. The method of Claim 12 wherein said solvent is
5 ethanol.

18. The method of Claim 12 wherein said water soluble
carrier is polyethylene glycol (PEG).

10 19. A method of treating an HIV infection comprising
administering an effective amount of a solid dispersion of
Claim 1 to a mammal in need of such treatment.

20. The method of Claim 19 wherein said HIV protease
15 inhibitor is selected from the group consisting of
(2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-isopropyl-4-
thiazolyl)methyl)amino)carbonyl)L-valinyl)amino-2-(N-((5-
thiazolyl)methoxy-carbonyl)-amino)-amino-1,6-diphenyl-3-
hydroxyhexane (ritonavir) and (2S,3S,5S)-2-(2,6-
20 Dimethylphenoxyacetyl)amino-3-hydroxy-5-[2S-(1-tetrahydro-
pyrimid-2-onyl)-3-methyl butanoyl] amino-1,6-diphenylhexane
(ABT-378).

21. The method of Claim 19 wherein said HIV protease inhibitor is a combination of (2S,3S,5S)-5-(N-(N-(N-methyl-N-((2-isopropyl-4-thiazolyl)methyl)amino)carbonyl)L-valinyl)amino-2-(N-((5-thiazolyl)methoxy-carbonyl)-amino)-amino-1,6-diphenyl-3-hydroxyhexane (ritonavir) and (2S,3S,5S)-2-(2,6-Dimethylphenoxyacetyl)amino-3-hydroxy-5-[2S-(1-tetrahydro-pyrimid-2-onyl)-3-methyl butanoyl] amino-1,6-diphenylhexane (ABT-378).

**Figure 1**

Amorphous ABT-538 Dispersions in PEG 8000
(Crossover Design, n=7)





Figure 2

**Figure 3**

**ABT-378 Dissolution in 0.1 N HCl at 37°C
(3.6 mg drug per capsule)**





Figure 4

**Figure 5**

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 May 2001 (17.05.2001)

PCT

(10) International Publication Number
WO 01/34118 A3

(51) International Patent Classification⁷: **A61K 9/14**,
31/427, 31/513, A61P 31/18

(21) International Application Number: PCT/US00/30910

(22) International Filing Date:
10 November 2000 (10.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/439,514 12 November 1999 (12.11.1999) US

(71) Applicant: **ABBOTT LABORATORIES** [US/US]; Dept.
377, Bldg. AP6D-2, 100 Abbott Park Road, Abbott Park,
IL 60064-6050 (US).

(72) Inventors: **FORT, James, J.**; 2700 Leaffield Terrace, Mid-
lothian, VA 23113 (US). **KRILL, Steven, L.**; 44 Mey-
ersville Road, Chatham, NJ 07928 (US). **LAW, Devalina**;

835C Country Club Drive, Libertyville, IL 60048 (US).
QIU, Yihong; 6118 Honeysuckle Lane, Gurnee, IL 60031
(US). **PORTER, William, R.**; 20 Manchester Lane, Ver-
non Hills, IL 60061 (US). **SCHMITT, Eric, A.**; 319 Ever-
green Court, Libertyville, IL 60048 (US).

(74) Agents: **CROWLEY, Steven, R.** et al.; Dept. 377 Bldg.
AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-
6050 (US).

(81) Designated States (*national*): CA, JP, MX.

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).


Published:
— with international search report

(88) Date of publication of the international search report:
10 January 2002

[Continued on next page]

(54) Title: SOLID DISPERSION PHARMACEUTICAL FORMULATIONS

Amorphous ABT-538 Dispersions in PEG 8000
Dissolution in 0.1N HCl at 37°C



(57) Abstract: A pharmaceutical composition is disclosed which comprises a solid dispersion of an HIV protease inhibitor in a water soluble carrier, such as PEG, having enhanced bioavailability and improved dissolution properties. The solid dispersion may optionally be encapsulated in hard gelatin capsules, compressed into a tablet, or may be granulated with a pharmaceutically acceptable granulating agent. Also disclosed are methods of making said solid dispersion and methods of treating an HIV infection employing said solid dispersion.

WO 01/34118 A3



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/L 00/30910

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K9/14 A61K31/427 A61K31/513 A61P31/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	B.J. AUNGST ET AL.: "Amphiphilic vehicles improve the oral bioavailability of a poorly soluble HIV protease inhibitor at high doses" INTERNATIONAL JOURNAL OF PHARMACEUTICS, vol. 156, no. 1, 1997, pages 79-88, XP001009732 Amsterdam (NL) cited in the application page 81, paragraph 2.2 page 84; table 2	1-5,9,19
A	---	6-8, 10-18, 20,21
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

19 June 2001

Date of mailing of the international search report

04/07/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Benz, K

INTERNATIONAL SEARCH REPORT

International Application No

PCT/ 00/30910

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 07696 A (ABBOTT LABORATORIES) 23 March 1995 (1995-03-23) page 1, paragraph 2 page 6, paragraph 4 page 17, paragraph 1 page 22; example 5 page 24; example 15 -----	1-21

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 00/30910

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9507696 A	23-03-1995	AU 695516 B	13-08-1998
		AU 7717694 A	03-04-1995
		CA 2167412 A	23-03-1995
		EP 0719142 A	03-07-1996
		IL 110752 A	26-07-2000
		JP 9502715 T	18-03-1997
		US 5948436 A	07-09-1999
		US 5725878 A	10-03-1998
		US 5876749 A	02-03-1999
